



PATIENT ASSESSED
REQUIRES BIOFILM- BASED WOUND CARE

PATIENT REQUIRES ANTISEPTIC DRESSING
FROM ACTIVHEAL ANTIMICROBIAL RANGE



USE FOR ALL AGES

OK FOR USE

CAN BE REMOVED

MAY REDUCE PAIN

SUITABLE FOR ALL AREAS
AND SACRAL SHAPED DRESSINGS

SUITABLE FOR SUPERFICIAL
AND NON-CAVITY WOUNDS

SUITABLE FOR USE
ON SLOUGHY WOUNDS

SUITABLE FOR LOCALISED INFECTION OR
SYSTEMIC INFECTION WITH ANTIBIOTICS

FOR USE ON MODERATE
TO HIGH EXUDATE WOUNDS

AGE
HOW OLD IS THE PATIENT?

SEX
IS THE PATIENT PREGNANT?

CAPACITY
CAPACITY OF THE PATIENT

WOUND PAIN
IS THE PATIENT SHOWING SIGNS OF
PAIN CAUSED BY THE WOUND?

WOUND LOCATION
IS THE WOUND AREA
HARD TO DRESS?

WOUND DEPTH
APPROPRIATE FOR USE
ON CAVITY WOUND?

DEVITALISED TISSUE
IS THE WOUND DEVITALISED?

INFECTION
IS THE WOUND INFECTED?

EXUDATE
HOW MUCH EXUDATE DOES
THE WOUND HAVE?

CAUTION WITH INFANTS

CAUTION IF
BREAST FEEDING

CAN BE REMOVED

MAY CAUSE
TRANSIENT STINGING

SUITABLE FOR ALL AREAS

SUITABLE FOR SUPERFICIAL
AND CAVITY WOUNDS

MOST EFFECTIVE WHEN
WOUND BED IS CLEAN

SUITABLE FOR LOCALISED INFECTION OR
SYSTEMIC INFECTION WITH ANTIBIOTICS

FOR USE ON MODERATE
TO HIGH EXUDATE WOUNDS

CAUTION WITH INFANTS

CAUTION IF
BREAST FEEDING

CAN FACILITATE CONCORDANCE
WITH SILVER TO THE WOUND BED

MAY CAUSE
TRANSIENT STINGING

SUITABLE FOR ALL AREAS
INCLUDING HARD TO DRESS AREAS

SUITABLE FOR SUPERFICIAL
AND CAVITY WOUNDS

MOST EFFECTIVE WHEN
WOUND BED IS CLEAN

SUITABLE FOR LOCALISED INFECTION OR
SYSTEMIC INFECTION WITH ANTIBIOTICS

FOR USE WITH LOW AND
MODERATE EXUDATE.

ABSORPTION CAN BE INCREASED BY CHOICE OF SECONDARY DRESSING.

IF THERE IS ANY KNOWN OR SUSPECTED SENSITIVITY TO THE ANTIMICROBIAL AGENT
OR THE CARRIER DRESSING THE PRODUCT SHOULD NOT BE USED.