



**PATIENT ASSESSED**  
REQUIRES BIOFILM- BASED WOUND CARE

**PATIENT REQUIRES ANTISEPTIC DRESSING**  
FROM ACTIVHEAL ANTIMICROBIAL RANGE



USE FOR ALL AGES

OK FOR USE

CAN BE REMOVED

MAY REDUCE PAIN

SUITABLE FOR ALL AREAS  
AND SACRAL SHAPED DRESSINGS

SUITABLE FOR SUPERFICIAL  
AND NON-CAVITY WOUNDS

SUITABLE FOR USE  
ON SLOUGHY WOUNDS

SUITABLE FOR LOCALISED INFECTION OR  
SYSTEMIC INFECTION WITH ANTIBIOTICS

FOR USE ON MODERATE  
TO HIGH EXUDATE WOUNDS

**AGE**  
HOW OLD IS THE PATIENT?

**SEX**  
IS THE PATIENT PREGNANT?

**CAPACITY**  
CAPACITY OF THE PATIENT

**WOUND PAIN**  
IS THE PATIENT SHOWING SIGNS OF  
PAIN CAUSED BY THE WOUND?

**WOUND LOCATION**  
IS THE WOUND AREA  
HARD TO DRESS?

**WOUND DEPTH**  
APPROPRIATE FOR USE  
ON CAVITY WOUND?

**DEVITALISED TISSUE**  
IS THE WOUND DEVITALISED?

**INFECTION**  
IS THE WOUND INFECTED?

**EXUDATE**  
HOW MUCH EXUDATE DOES  
THE WOUND HAVE?

CAUTION WITH INFANTS

CAUTION IF  
BREAST FEEDING

CAN BE REMOVED

MAY CAUSE  
TRANSIENT STINGING

SUITABLE FOR ALL AREAS

SUITABLE FOR SUPERFICIAL  
AND CAVITY WOUNDS

MOST EFFECTIVE WHEN  
WOUND BED IS CLEAN

SUITABLE FOR LOCALISED INFECTION OR  
SYSTEMIC INFECTION WITH ANTIBIOTICS

FOR USE ON MODERATE  
TO HIGH EXUDATE WOUNDS

CAUTION WITH INFANTS

CAUTION IF  
BREAST FEEDING

CAN FACILITATE CONCORDANCE  
WITH SILVER TO THE WOUND BED

MAY CAUSE  
TRANSIENT STINGING

SUITABLE FOR ALL AREAS  
INCLUDING HARD TO DRESS AREAS

SUITABLE FOR SUPERFICIAL  
AND CAVITY WOUNDS

MOST EFFECTIVE WHEN  
WOUND BED IS CLEAN

SUITABLE FOR LOCALISED INFECTION OR  
SYSTEMIC INFECTION WITH ANTIBIOTICS

FOR USE WITH LOW AND  
MODERATE EXUDATE.

ABSORPTION CAN BE INCREASED BY CHOICE OF SECONDARY DRESSING.

IF THERE IS ANY KNOWN OR SUSPECTED SENSITIVITY TO THE ANTIMICROBIAL AGENT  
OR THE CARRIER DRESSING THE PRODUCT SHOULD NOT BE USED.